NCBI Map Viewer Tutorial

NCBI's Map Viewer can be used to visualize an organism's genome and annotation. The organisms represented in the Map Viewer include human, mouse, rat, zebrafish, mosquito, nematode, fruit fly, yeast, arabidopsis and others.

In these exercises, we will use the human genome Map Viewer to

- Locate a human gene
- Download the gene sequence along with its upstream sequence (to analyze promoter regions)
- Determine whether the sequence is in the draft or finished form
- Find a possible splice variant using Model Maker
- Identify whether variations in the gene are associated with a disease
- Obtain information about the mouse and rat homologs
- Obtain phenotype information

The first exercise includes screenshots of the suggested steps as well as a few introductory slides.

Exercise 1

We will use as an example the glial cell derived neurotrophic factor GDNF. GDNF may contribute to Hirschsprung disease when mutated. Search for "GDNF" in the Map Viewer. Access the Human Map Viewer page by clicking on the Map Viewer link in the footer of the NCBI home page, then selecting the "Homo sapiens (human) Build 37.1" link. Enter GDNF as a search term and click on the "Find" button. Filter your search result for gene maps by checking the "Gene" box in the "Quick Filter" menu.

Name the chromosome on which this gene is located. Click on the Genes_seq map link of the map element GDNF in the reference assembly. What is the map that is displayed? Turn on the ruler line for the map through the Maps & Options link. What are the nucleotide locations for the gene on the chromosome? What is the orientation of the gene on the chromosome? Download the gene sequence by using the "dl" link. (Change the strand, if necessary). In order to search for promoter elements, you may also download 5000 nucleotides of upstream sequence by adjusting the locations for the upstream 5000 nucleotides.

Add the Clone, Component and Contig maps for this region. Name the contig and GenBank accession numbers for the sequence covering this region. Are the sequences in the finished form? Is there a clone mapped to this region? If so, how can you order it?

Remove all the maps except Genes_seq and add the Ab initio (model) and RefSeq Transcripts maps. Does the gene prediction match the current gene annotation? How many alternatively spliced transcripts have been annotated for the gene? Display the current data as "Data As Table View".

Using the Model Maker (mm), obtain a possible alternatively spliced product and its translated amino acid sequence. Search for similar proteins by using BLAST.

Remove all the maps except Genes_seq and add the Gene maps for mouse, chimp and rat. Are the gene structures in the three organisms similar?


Remove all the maps except the human Gene map, and add the phenotype map. Name the disease with which the GDNF gene is associated. Obtain more information about the disease by linking to the corresponding OMIM record.



What is Map Viewer?

Map Viewer allows you to view and search an organism's complete genome, display chromosome maps, and zoom into progressively greater levels of detail, down to the sequence data for a region of interest.

It allows for the analysis of genes and genome annotation in graphical format.






- · Locate a human gene
- Download the gene sequence along with its upstream region (for promoter analysis)
- Determine whether the sequence is in draft or finished form
- Find a possible splice variant via Model Maker
- Identify whether variations in the gene are associated with a disease
- Obtain information about mouse and rat homologues

High Throughput Genomic Sequences


Sequencing of BACs:

Draft:

Phase 0 – single/few pass reads of a single clone Phase 1 – unfinished, unordered BACs with gaps




Phase 2 - unfinished, ordered with gaps





Finished:


Phase 3 - high quality finished sequence with no gaps































Homo sapiens (human) (Build 37.1)
Region to retrieve (in chromosome coordinates):
Chromosome: 5 Strand: plus \$
from: 37815753 adjust by: -0K
to: 37839782 adjust by: +0K Change Region/Strand
Sequence Format: FASTA :
This chromosome region corresponds to the contig region(s):
Contig start stop strand
NT_006576.16 37805753 37829782 + <u>Display</u> <u>Save to Disk</u> <u>View Evidence</u> <u>ModelMaker</u>
Homo sapiens (human) (Build 37.1)
Region to retrieve (in chromosome coordinates):
Region to retrieve (in chromosome coordinates):
Region to retrieve (in chromosome coordinates): Chromosome: 5 Strand: minus \$
Region to retrieve (in chromosome coordinates): Chromosome: 5 Strand: minus \$ from: 37815753 adjust by: -0K
Region to retrieve (in chromosome coordinates): Chromosome: 5
















monto sap	<i>piens</i> (hur	man) Build 37	1.1 (0	Current)							BLAST The I	Human Genome
Data As Tal	ble View							<u>Do</u>	wnload All			
Genes On S	Sequence				All S	Sequence N	<u>Iaps</u>					next
	on Chron	,798K-37,858K t nosome: 1536	bp						Downlo	oad/View Sequence/Eviden	nce	Download Date
	stop	Symbol	0			Links		<u>E</u>	Cyto	Description		
37813681	37814281	LOC100287032	-		sv	dl ev mm		mRNA	5	similar to hCG2006445		
37815753	37839782	<u>GDNF</u>	- <u>O</u>	MIM HGNO	C sv pr	dl ev mm h	m sts CCDS S	NP best RefSeq	5p13.1-p12	glial cell derived neurotro	ophic factor	
Ab initio me	odels			All S	equen	ce Maps					prev	next
	ls On Chro	,798K-37,858K tomosome: 3410	р						Downlo	oad/View Sequence/Eviden	nce	Download Dat
start	stop	Symbol	O Be	est hit	Best l	hit descript	ion					
37805733	37812363	hmm1374634	-									
37813681	37814281	hmm1376634	-									
				P 000505.1	glial	cell derived	I neurotrophic	factor isoform	1 preproprot	ein [Homo sapiens]		
37815753	37835929	hmm12438633	- <u>N</u> I							ein [Homo sapiens] ein [Homo sapiens]		
37815753 37815753	37835929 37839776	hmm12438633	- <u>NI</u> - <u>NI</u>									
37815753 37815753 37836700 RefSeq Trai	37835929 37839776 37845817 nscripts Or	hmm12438633 hmm12436633 hmm1378634	- <u>NI</u> - <u>NI</u> +				I neurotrophic		1 preproprot	ein [Homo sapiens]	prev	
37815753 37815753 37836700 RefSeq Tran Region Disp	37835929 37839776 37845817 nscripts Or played: 37 , eq Transcrip	hmm12438633 hmm12436633 hmm1378634 in Sequence 798K-37,858K it pts On Chromoso	- <u>NF</u> - <u>NF</u> +	P 000505.1			I neurotrophic	factor isoform	1 preproprot			Download Da
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran	37835929 37839776 37845817 nscripts Or played: 37 , eq Transcrip	hmm12438633 hmm12436633 hmm1378634 in Sequence 798K-37,858K it pts On Chromoso	- NF - NF +	P 000505.1		cell derived	I neurotrophic	factor isoform	1 preproprot	ein [Homo sapiens]		
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran start	37835929 37839776 37845817 nscripts Or played: 37, eq Transcripts in stop	hmm12438633 hmm12436633 hmm1378634 in Sequence 798K-37,858K it pts On Chromoso Region: 4	- NF - NF +	P 000505.1 1527 ocus	glial	cell derived	All Sequinks	factor isoform ence Maps Align quality 1	1 preproprot	ein [Homo sapiens]		
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran start 37813681	37835929 37839776 37845817 nscripts Or played: 37 , eq Transcripts in stop 37814281	hmm12438633 hmm12436633 hmm1378634 n Sequence 798K-37,858K b pts On Chromoso Region: 4 Accession	- NF - NF + bp ome::	P 000505.1 1527 ocus	<u>O</u>	cell derived	All Sequinks BLink	ence Maps Align quality 1 identical	Downlor Description Similar to hCo	ein [Homo sapiens]		
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran start 37813681	37835929 37839776 37845817 Inscripts Or played: 37 , eq Transcripts in stop 37814281 37834898	hmm12438633 hmm12436633 hmm1378634 1 Sequence 798K-37,858K ly 198 On Chromoso Region: 4 Accession XM 002342491 NM 199234.1	- NF - NF + bp ome::	P 000505.1 1527 ocus OC1002870	<u>O</u>	L sv prev	All Sequinks BLink	factor isoform ence Maps Align quality 1 identical sidentical sidentical s	Downlor Description Similar to hCo	ein [Homo sapiens] pad/View Sequence/Eviden G2006445		
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran start 37813681 37815753	37835929 37839776 37845817 nscripts Or played: 37, eq Transcript stop 37814281 37834898 37816100	hmm12438633 hmm12436633 hmm1378634 1 Sequence 798K-37,858K ly 198 On Chromoso Region: 4 Accession XM 002342491 NM 199234.1 exon	- NF - NF + bp ome::	P 000505.1 1527 ocus OC1002870	<u>O</u>	L sv prev	All Sequinks BLink	Align quality I identical sidentical gCDS 3	Downlo	ein [Homo sapiens] pad/View Sequence/Eviden G2006445		
37815753 37815753 37836700 RefSeq Tran Region Disp Total RefSe RefSeq Tran start 37813681 37815753 37815753	37835929 37839776 37845817 nscripts Or played: 37, aq Transcript stop 37814281 37834898 37816100 37834836	hmm12438633 hmm12436633 hmm1378634 1 Sequence 7798K-37,858K ly 15 On Chromoso Region: 4 Accession XM 002342491 NM 199234.1 exon intron	- NF - NF + bp ome::	P 000505.1 1527 ocus OC1002870	<u>O</u>	L sv prev	All Sequinks BLink	Align quality I identical sidentical gCDS	Downless Description Similar to hCoglial cell derivates by	ein [Homo sapiens] pad/View Sequence/Eviden G2006445		
37815753 37836700 RefSeq Tran Region Disj Total RefSe RefSeq Tran start 37813681 37815753 37815753 37816101	37835929 37839776 37845817 nscripts Or played: 37, q Transcrip nscripts in stop 37814281 37834898 37816100 37834836 37834890	hmm12438633 hmm12436633 hmm1378634 in Sequence 7/98K-37,858K bpts On Chromoso Region: 4 Accession XM 003342491 NM 199234.1 exon intron exon	- NF - NF + bp ome::	P 000505.1 1527 ocus OC1002870	<u>O</u>	L sv prev	All Sequinks BLink	factor isoform ence Maps Align quality I identical sidentical gCDS CDS CDS	Downlos Description similar to hCoglial cell derivates bp 18736 bp	ein [Homo sapiens] pad/View Sequence/Eviden G2006445		
37815753 37815753 37836700 RefSeq Tran Region Dis Total RefSe RefSeq Tran start 37813681 37815753 37815753 37816101 37834837 37834891	37835929 37839776 37845817 nscripts Or played: 37, pq Transcript in stop 37814281 37834898 37816100 37834896 37834890 37834898	hmm12438633 hmm12436633 hmm1378634 in Sequence 7/98K-37,858K bpts On Chromoso Region: 4 Accession XM 003342491 NM 199234.1 exon intron exon	- NE + + LC G	P 000505.1 1527 ocus OC1002870	O 32 - - - -	L syprey ug sy prey	All Sequinks BLink	factor isoform ence Maps Align quality I identical identical graph (CDS CDS CDS CDS CDS CDS CDS CDS CDS CDS	Description similar to hCt glial cell derivates by 18736 bp 54 bp 8 bp	ein [Homo sapiens] pad/View Sequence/Eviden G2006445		
37815753 37815753 37836700 RefSeq Tran Region Dis Total RefSe RefSeq Tran start 37813681 37815753 37815753 37816101 37834837 37834891	37835929 37839776 37845817 nscripts Or played: 37, eq Transcripts in stop 37814281 37834898 37816100 37834896 37834898 37834898 37835593	hmm12438633 hmm12436633 hmm1378634 in Sequence 798K-37,858K t pts On Chromoso Region: 4 Accession XM 002342491 NM 199234.1 exon intron exon exon	- NE + + LC G	P 000505.1 1527 ocus OC1002870	O 32 - - - -	L syprey ug sy prey	All Sequinks BLink BLink	factor isoform ence Maps Align quality I identical identical graph (CDS CDS CDS CDS CDS CDS CDS CDS CDS CDS	Description similar to hCt glial cell derivates by 18736 bp 54 bp 8 bp	ein [Homo sapiens] pad/View Sequence/Eviden G2006445 ved neurotrophic factor		





















Alignments >ref|XP 001143054.1| PREDICTED: similar to neurotrophic factor isoform 3 [Pan troglodytes] gb|EAW55964.1| glial cell derived neurotrophic factor, isoform CRA b [Homo sapiens] Length=228 Score = 432 bits (1112), Expect = 4e-120 Identities = 211/211 (100%), Positives = 211/211 (100%), Gaps = 0/211 (0%) MKLWDVVAVCLVLLHTASAFPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQ 60 Query 1 MKLWDVVAVCLVLLHTASAFPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQ Sbjct 18 MKLWDVVAVCLVLLHTASAFPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQ 77 Query 61 FDDVMDFIOATIKRLKRSPDKOMAVLPRRERNROAAAANPENSRGKGRRGORGKNRGCVL 120 ${\tt FDDVMDFIQATIKRLKRSPDKQMAVLPRRERNRQAAAANPENSRGKGRRGQRGKNRGCVL}$ FDDVMDFIQATIKRLKRSPDKQMAVLPRRERNRQAAAANPENSRGKGRRGQRGKNRGCVL 137 Sbjct 78 Query 121 TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR 180 TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR Sbjct 138 TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR 197 Query 181 PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI 211 PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI Sbjct 198 PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI 228

```
>qb|AAX36770.1| qlial cell derived neurotrophic factor [synthetic construct]
Length=212
Score = 432 bits (1112), Expect = 7e-120
Identities = 211/211 (100%), Positives = 211/211 (100%), Gaps = 0/211 (0%)
Query 1
           MKLWDVVAVCLVLLHTASAFPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQ
           MKLWDVVAVCLVLLHTASAF PLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDO
Sbjct 1
           MKLWDVVAVCLVLLHTASAFPLPAGKRPPEAPAEDRSLGRRRAPFALSSDSNMPEDYPDQ
          FDDVMDFTOATTKRIKRSPDKOMAVI,PRRERNROAAAANPENSRGKGRRGORGKNRGCVI. 120
Query 61
           FDDVMDFIQATIKRLKRSPDKQMAVLPRRERNRQAAAANPENSRGKGRRGQRGKNRGCVL
Sbjct 61
          FDDVMDFIQATIKRLKRSPDKQMAVLPRRERNRQAAAANPENSRGKGRRGQRGKNRGCVL 120
Query 121 TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR 180
           TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR
Sbjct 121 TAIHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCR 180
Query 181 PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI 211
           PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI
Sbjct 181 PIAFDDDLSFLDDNLVYHILRKHSAKRCGCI
>ref[NP_000505.1] glial cell derived neurotrophic factor isoform 1 preproprotein
[Homo sapiens]
ref | XP 526944.1 | PREDICTED: glial cell derived neurotrophic factor isoform 4 [Pan
roalodutoal
```









Exercise 2

Search for "PRNP" in the Map Viewer. Access the Human Map Viewer page by clicking on <u>Map Viewer</u>, then selecting the "Homo sapiens (human) Build 36" link. Enter PRNP as a search term.

Name the chromosome on which this gene is located. Click on the Genes_seq map link of the map element PRNP in the reference assembly.

What is the map that is displayed? Turn on the ruler line for the map through the Maps & Options link. What are the nucleotide locations for the gene on the chromosome? What is the orientation of the gene on the chromosome?

Download the gene sequence by using the "dl" link. (Change the strand, if necessary). In order to search for promoter elements, you may also download 5000 nucleotides of upstream sequence by adjusting the locations for the upstream 5000 nucleotides.

Add the Clone, Component and Contig maps for this region. Name the contig and GenBank accession numbers for the sequence covering this region. Are the sequences in the finished form? Is there a clone mapped to this region? If so, how can you order it?

Remove all the maps except Genes_seq and add the Ab initio (model) and RefSeq Transcripts maps. Does the gene prediction match the current gene annotation? How many alternatively spliced transcripts have been annotated for the gene? Display the current data as "Data As Table View".

Using the Model Maker (mm), obtain a possible alternatively spliced product and its translated amino acid sequence. Search for similar proteins by using BLAST.

Remove all the maps except Genes_seq and add the Gene maps for mouse, chimp and rat. Are the gene structures in the three organisms similar?

Remove all the maps except the human Gene map, and add the phenotype map. Name the disease with which the PRNP gene is associated. Obtain more information about the disease by linking to the corresponding OMIM record.